

## 5.4 Indefinite Integrals and the Net Change Theorem

An indefinite integral is one in which the limits of integration are not given. That is, an **indefinite integral** is

$$\int f(x) dx = F(x) \text{ means } F'(x) = f(x)$$

**Example 1** Find the general indefinite integral:  $\int(\cos(x) + x^2) dx$ .

### Table of Indefinite Integrals

Complete the table of indefinite integrals

- |                                      |  |
|--------------------------------------|--|
| 1. $\int c f(x) dx = c \int f(x) dx$ | 2. $\int [f(x) \pm g(x)] dx = \int f(x) dx + \int g(x) dx$ |
| 3. $\int x^n dx =$                   | 4. $\int \frac{1}{x} dx =$                                 |
| 5. $\int e^x dx =$                   | 6. $\int a^x dx =$   |
| 7. $\int \sin(x) dx =$               | 8. $\int \cos(x) dx =$                                     |
| 9. $\int \sec^2(x) dx =$             | 10. $\int \csc^2(x) dx =$                                  |
| 11. $\int \sec(x) \tan(x) dx =$      | 12. $\int \csc(x) \cot(x) dx =$                            |
| 13. $\int \frac{1}{x^2+1} dx =$      | 14. $\int \frac{1}{\sqrt{1-x^2}} dx$                       |

**Example 2** Find the general indefinite integral:  $\int \left( \frac{x^2+1}{x} + 2^x \right) dx$

**Example 3** Find the general indefinite integral:  $\int \frac{\sin(x)}{\cos^2(x)} dx$

**Example 4** Evaluate the definite integral and interpret the result:  $\int_1^4 (x^2 - 3x) dx$ .

Suppose the function  $f(x) = x^2 - 3x$  in example (4) represented the rate of growth in the value of a certain stock. The net area, or **net change**, represents the change in the value of the stock over that time interval. This is simply another way to reformulate **FTC2**:

---

### The Net Change Theorem

The integral of a rate of change is the net change:

$$\int_a^b f'(x) dx = F(b) - F(a)$$

---

**Example 5** The velocity of a particle moving on the number line is given by  $v(t) = t^2 - 4t + 3$ . Find its net change of position, or *displacement*, on the interval  $[1, 5]$ , and find the total distance traveled.

**Example 6** The rate the altitude of a jet is changing is given by  $A'(t) = t^3 - 6t$ . Find a value  $c > 1$  such that the net change on the interval  $[1, c]$  is zero.