

6.1-3 Orthogonality

Inner Product

Let $\mathbf{u} = \begin{pmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{pmatrix}$ and $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix}$. The **inner product** of \mathbf{u} and \mathbf{v} is $(u_1 \ u_2 \ \cdots \ u_n) \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} = u_1 v_1 + u_2 v_2 + \cdots + u_n v_n$. This is also called the **dot product** $\mathbf{u} \cdot \mathbf{v}$.

Length of a Vector

The length, or **norm**, of a vector is $\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$. Also note, $\|\mathbf{v}\|^2 = \mathbf{v} \cdot \mathbf{v}$.

A vector whose length is 1 is called a **unit vector** and is found by dividing by its norm: $\mathbf{u} = \frac{\mathbf{v}}{\|\mathbf{v}\|}$. The process of creating \mathbf{u} from \mathbf{v} is called **normalizing** \mathbf{v} .

Distance Between Two Vectors

For vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n , the **distance between \mathbf{u} and \mathbf{v}** , written as $\text{dist}(\mathbf{u}, \mathbf{v})$ is the length of the vector $\mathbf{u} - \mathbf{v}$, that is $\text{dist}(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\|$.

Example 1 Find the distance between the vectors $\mathbf{u} = \begin{pmatrix} 7 \\ 3 \end{pmatrix}$ and $\mathbf{v} = \begin{pmatrix} 2 \\ 4 \end{pmatrix}$, and draw a sketch of the three vectors \mathbf{u} , \mathbf{v} , and $\mathbf{u} - \mathbf{v}$.

Orthogonal Vectors

Recall from calculus that another definition of the dot product is $\mathbf{u} \cdot \mathbf{v} = \|\mathbf{u}\| \|\mathbf{v}\| \cos(\theta)$ where θ is the angle between \mathbf{u} and \mathbf{v} . If the vectors are in \mathbb{R}^2 or \mathbb{R}^3 and are perpendicular, then $\theta = 90^\circ$. This means $\mathbf{u} \cdot \mathbf{v} = 0$. In general, two vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n are **orthogonal** if $\mathbf{u} \cdot \mathbf{v} = 0$.

Example 2 Determine which vectors are orthogonal: $\mathbf{u}_1 = \begin{pmatrix} 2 \\ 3 \\ -4 \end{pmatrix}$, $\mathbf{u}_2 = \begin{pmatrix} -2 \\ 1 \\ 1 \end{pmatrix}$, $\mathbf{u}_3 = \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$.

Orthogonal Complement

Let W be a subspace of \mathbb{R}^n . If \mathbf{z} is orthogonal to every vector in W then \mathbf{z} is said to be orthogonal to W . The set of all vectors \mathbf{z} that are orthogonal to W is called the **orthogonal complement** of W and is denoted by W^\perp . This leads to two properties

- (1) A vector \mathbf{x} is in W^\perp if and only if \mathbf{x} is orthogonal to every vector in a set that spans W .
- (2) W^\perp is a subspace of \mathbb{R}^n .

Note: For an $m \times n$ matrix A $(\text{Row } A)^\perp = \text{Nul } A$ and $(\text{Col } A)^\perp = \text{Nul } A^T$. (Thm 6.1.3)

Orthogonal Sets

A set of vectors $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n\}$ is said to be an orthogonal set if each pair of vectors $\mathbf{u}_i \cdot \mathbf{u}_j = 0$ for all $i \neq j$.

Example 3 Show that $\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$ is an orthogonal set, where, $\mathbf{u}_1 = \begin{pmatrix} 2 \\ -1 \\ 0 \end{pmatrix}$, $\mathbf{u}_2 = \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$ and $\mathbf{u}_3 = \begin{pmatrix} -2 \\ -4 \\ 5 \end{pmatrix}$.

Orthogonal Basis

An **orthogonal basis** for a subspace W of \mathbb{R}^n is a basis for W that is also an orthogonal set.

Theorem

Let $\{u_1, u_2, \dots, u_p\}$ be an orthogonal basis for a subspace W of \mathbb{R}^n . For each \mathbf{y} in W , the weights in the linear combination $\mathbf{y} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + \dots + c_p \mathbf{u}_p$ are given by $c_j = \frac{\mathbf{y} \cdot \mathbf{u}_j}{\mathbf{u}_j \cdot \mathbf{u}_j}$, ($j = 1, \dots, p$).

Example 4 The set $S = \{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$ of vectors in example 3 form an orthogonal basis for \mathbb{R}^3 . Express the vector $\mathbf{y} = \begin{pmatrix} 7 \\ 4 \\ -3 \end{pmatrix}$ in terms of S .

Orthonormal Sets; Orthonormal Basis

An orthogonal set of vectors $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n\}$ each of unit length is called an **orthonormal set**. If W is a subspace spanned by an orthonormal set, then $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n\}$ is an orthonormal basis.

Example 5 Give an example of two well known orthonormal bases.

Theorem An $m \times n$ matrix U has orthonormal columns if and only if $U^T U = I$.

Orthogonal Projection

Recall from calculus the quantity $c = \frac{\mathbf{y} \cdot \mathbf{u}}{\|\mathbf{u}\|}$ is the scalar projection of \mathbf{y} onto \mathbf{u} , or the amount of \mathbf{y} in the direction of \mathbf{u} .

The **orthogonal projection** is the vector $\hat{\mathbf{y}} = c \mathbf{u}$, or $\hat{\mathbf{y}} = \frac{\mathbf{y} \cdot \mathbf{u}}{\mathbf{u} \cdot \mathbf{u}} \mathbf{u}$. If L is the subspace spanned by \mathbf{u} then $\hat{\mathbf{y}} = \text{proj}_L \mathbf{u}$.

Example 6 Find the closest approximation to $\mathbf{y} = \langle 3, 1, 5, 1 \rangle$ spanned by $\mathbf{v}_1 = \langle 3, 1, -1, 1 \rangle$ and $\mathbf{v}_2 = \langle 1, -1, 1, -1 \rangle$